

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification ⁶ : C07H 17/08, A61K 31/70</p>	<p>A1</p>	<p>(11) International Publication Number: WO 96/18633 (43) International Publication Date: 20 June 1996 (20.06.96)</p>
<p>(21) International Application Number: PCT/EP95/04815 (22) International Filing Date: 7 December 1995 (07.12.95) (30) Priority Data: MI94A002496 13 December 1994 (13.12.94) IT (71) Applicant (for all designated States except US): ZAMBON GROUP S.P.A. [IT/IT]; Via della Chimica, 9, I-36100 Vicenza (IT). (72) Inventors; and (75) Inventors/Applicants (for US only): PELLACINI, Franco [IT/IT]; Via F.lli Rizzardi, 16/14, I-20151 Milano (IT). SCHIOPPACASSI, Giovanna [IT/IT]; Via Togliatti, 43, I-20017 Rho (IT). ALBINI, Enrico [IT/IT]; Via Torchietto, 14, I-27100 Pavia (IT). BOTTA, Daniela [IT/IT]; Via Valleggio, 4, I-22100 Como (IT). ROMAGNANO, Stefano [IT/IT]; Via D. Scarlatti, 33, I-20090 Buccinasco (IT). SANTANGELO, Francesco [IT/IT]; Via Don Gnocchi, 33, I-20148 Milano (IT).</p>		<p>(81) Designated States: AL, AM, AU, BB, BG, BR, BY, CA, CN, CZ, EE, FI, GE, HU, IS, JP, KG, KP, KR, KZ, LK, LR, LT, LV, MD, MG, MK, MN, MX, NO, NZ, PL, RO, RU, SG, SI, SK, TJ, TM, TT, UA, US, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG), ARIPO patent (KE, LS, MW, SD, SZ, UG). Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i></p>
<p>(54) Title: ERYTHROMYCIN A 9-O-OXIME DERIVATIVES ENDOWED WITH ANTIBIOTIC ACTIVITY</p> <p>(57) Abstract</p> <p>Compounds of formula (I), wherein A, R₁, R₂, M, n, m and r have the meanings reported in the description, processes for their preparation and pharmaceutical compositions containing them as active ingredients are described. The compounds of formula (I) are useful in the treatment of infectious diseases.</p> <div style="text-align: center;"> $\text{A}-(\text{CH}_2)_n-\overset{\text{R}_1}{\underset{ }{\text{N}}}-(\text{CH}_2)_m-\overset{\text{R}_2}{\underset{ }{\text{N}}}-(\text{CH}_2)_r-\text{O}-\text{N}=\text{M} \quad (\text{I})$ </div>		

FOR THE PURPOSES OF INFORMATION ONLY


Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Larvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

"Erythromycin A 9-O-oxime derivatives endowed with antibiotic activity"

5 The present invention relates to Erythromycin A derivatives endowed with antibiotic activity, useful in the treatment of infectious diseases and, more particularly, it relates to Erythromycin A 9-[O-(aminoalkyl)oxime] derivatives endowed with antibiotic activity against Gram-positive and Gram-negative microorganisms.

10 Erythromycin A [The Merck Index, XI Ed., No. 3626] is a well-known naturally occurring macrolide endowed with antibiotic activity, having the following structure



25

30

Besides being active against some non-bacterial microorganisms such as rickettsiae and mycoplasmas, Erythromycin A is endowed with antibacterial activity mainly against Gram-positive microorganisms such as streptococci, staphylococci and pneumococci, but it results effective as well against some Gram-negative microorganisms such as, for instance, Haemophilus influenzae, Neisseria gonorrhoeae and Bordetella pertussis.

In addition to the well-known activity against the aforementioned prokaryotes, it has been recently described in the literature that Erythromycin A and other macrolide antibiotics are active against eukaryotic parasites [P. A. Lartey et al., Advances in Pharmacology, 28, 307-343 (1994)].

Also in the case of Erythromycin A, likewise other antibacterial drugs, phenomena of resistance were observed with some bacterial strains.

In particular, the phenomenon was observed in the treatment of infections caused by staphylococci following to prolonged administration of Erythromycin A [A. Kucers and N. McK. Bennett, The use of antibiotics, A Comprehensive Review with Clinical Emphasis, William Heinemann Medical, IV Ed., (1987) 851-882].

The interest towards macrolide antibiotics derives from their use in clinical and veterinary therapy in the treatment of several infectious diseases such as, for instance, the infections of the respiratory tract, of the gastrointestinal tract, of the urogenital tract and of the external organs like skin, eye and ear.


Because of its high instability in acidic environment Erythromycin A is irreversibly converted, for instance in the gastrointestinal tract following to oral administration, into derivatives devoid of antibiotic activity, conferring thus poor bioavailability on the

active principle [H. A. Kirst, Annual Reports in Medicinal Chemistry, 25, 119-128 (1989)].

In order to overcome the above drawbacks, the research was addressed
5 to compounds which, while maintaining the good antibiotic properties of Erythromycin A, resulted to be characterized by a superior stability to the acids and better pharmacokinetic properties such as, for instance, superior oral bioavailability, haematic concentration, tissue penetration and half-life.

10 Within this field, we can cite as an example the carbamates and carbonates of Erythromycin A or Erythromycin A 9-O-oxime described in the European patent applications No. 0216169 and No. 0284203 (both in the name of Beecham Group PLC) and the compounds described in the European patent application No. 0033255 (Roussel-Uclaf).

15 The European patent application No. 0033255, in particular, describes derivatives of Erythromycin A 9-O-oxime of formula



wherein

Ery represents the Erythromycin A residue wherein the oxime group
20 (-N=Ery) is in place of the carbonyl group in position 9 (O=Ery); A represents a straight or branched alkyl group with from 1 to 6 carbon atoms; R represents, inter alia, an optionally substituted alkoxy group with from 1 to 6 carbon atoms, or a group [-N(R₁)R₂] wherein R₁ and R₂, the same or different, represent a hydrogen atom
25 or an optionally substituted alkyl group with from 1 to 6 carbon atoms.

The compounds described in the European patent application No. 0033255 such as, for instance, Erythromycin A 9-[O-[(2-methoxyethoxy)methyl]oxime], known with the International Nonproprietary Name Roxithromycin [The Merck Index, XI Ed., No. 8253], Erythromycin A


- 4 -

9-[O-[(2-dimethylamino)ethyl]oxime] and Erythromycin A
 9-[O-[(2-diethylamino)ethyl]oxime] have a spectrum of activity in
vitro comparable to that of Erythromycin A but are endowed with a
 5 superior stability to the acids and better pharmacokinetic
 properties.

Said compounds, therefore, present an antibiotic activity against
 Gram-positive bacteria such as staphylococci, streptococci and
 pneumococci and against some Gram-negative bacteria such as, for
 instance, Haemophilus influenzae and Haemophilus pertussis.

10 Now we have found compounds derivative of Erythromycin A 9-O-oxime
 and, more particularly, compounds derivative of Erythromycin A
 9-[O-(aminoalkyl)oxime], partly comprised but not exemplified in the
 European patent application No. 0033255, which have a wider spectrum
 of antibacterial activity against Gram-positive microorganisms and,
 15 particularly, against Gram-negative microorganisms, and improved
 pharmacokinetic properties such as, for instance, a superior dura-
 tion of action and a superior half-life of tissue elimination, with
 respect to the compounds described in the aforementioned European
 patent application.

20 Object of the present invention, therefore, are the compounds of
 formula



(I)

25 wherein

A is a phenyl group or a heterocycle with 5 or 6 members containing
 1 or more heteroatoms selected among nitrogen, oxygen and sul-
 phur, optionally substituted with from 1 to 3 groups, the same or
 different, selected among straight or branched C₁-C₄ alkyl or

30

alkoxy groups, C₁-C₂ alkylenedioxy groups, C₁-C₄ alkylsulphonyl groups, phenyl, phenoxy, hydroxy, carboxy, nitro, halogen and trifluoromethyl groups;


5 R₁ and R₂, the same or different, represent a hydrogen atom or a straight or branched C₁-C₄ alkyl groups;

n is 1 or 2;

m is an integer comprised between 1 and 8;

p is an integer comprised between 2 and 6;

10 M represents a group of formula



25 wherein

R₃ is a hydrogen atom or a methyl group;

and pharmaceutically acceptable salts thereof.

The oximes of formula (I) can have Z or E configuration.

Object of the present invention, therefore, are the compounds of formula (I) having Z or E configuration, with a preference for these

30

latter.

The compounds of formula (I) are endowed with antibiotic activity and are characterized by a high stability to the acids and by good pharmacokinetic properties, being thus used in human or veterinary therapy for the treatment of several infectious diseases such as, for instance, the infections of the central nervous system, of the upper and lower respiratory tract, of the gastrointestinal tract, of the urogenital tract, of the odontological tissue and of the external organs such as skin, eye and ear.

10 In the present description, unless otherwise specified, with the term C₁-C₄ alkyl group we intend a straight or branched C₁-C₄ alkyl such as methyl, ethyl, n.propyl, isopropyl, n.butyl, isobutyl, sec-butyl and tert-butyl group; with the term C₁-C₄ alkoxy group we intend a straight or branched C₁-C₄ alkoxy such as methoxy, ethoxy, 15 n.propoxy, isopropoxy, n.butoxy, isobutoxy, sec-butoxy and tert-butoxy group; with the term C₁-C₂ alkylenedioxy group we intend a methylenedioxy or ethylenedioxy group.

20 With the term heterocycle with 5 or 6 members containing 1 or more heteroatoms selected among nitrogen, oxygen and sulphur we intend a heterocycle preferably selected among pyridine, pyrrole, pyrrolidine, furan, tetrahydrofuran and thiophene.

Preferred compounds are the compounds of formula (I) wherein A represents a phenyl group or a heterocycle selected between pyridine and furan, optionally substituted with from 1 to 3 groups selected 25 among hydroxy, methoxy, methylenedioxy, n.butoxy, phenoxy, phenyl, methylsulphonyl, nitro, halogen and trifluoromethyl groups; R₁ and R₂, being the same, represent a hydrogen atom or a methyl group; R₃ represents a hydrogen atom.

Still more preferred compounds are the compounds of formula (I)

wherein A represents a phenyl group optionally substituted with a group selected among phenoxy, nitro and trifluoromethyl; R₁ and R₂, being the same, represent a hydrogen atom or a methyl group; n is equal to 1; m is equal to 6; r is equal to 2; R₃ represents a hydrogen atom.

Pharmaceutically acceptable salts of the compounds of formula (I) are the salts with organic or inorganic acids such as, for instance, hydrochloric, hydrobromic, hydroiodic, nitric, sulphuric, phosphoric, acetic, tartaric, citric, benzoic, succinic and glutaric acid.

Specific examples of preferred compounds of formula (I) are:

Erythromycin A (E)-9-[0-[2-[6-(phenylmethylamino)hexylamino]ethyl]-oxime];

Erythromycin A (E)-9-[0-[2-[2-(phenylmethylamino)ethylamino]ethyl]-oxime];

Erythromycin A (E)-9-[0-[6-[6-(phenylmethylamino)hexylamino]hexyl]-oxime];

Erythromycin A (E)-9-[0-[6-[3-(phenylmethylamino)propylamino]hexyl]-oxime];

Erythromycin A (E)-9-[0-[6-[5-(phenylmethylamino)pentylamino]hexyl]-oxime];

Erythromycin A (E)-9-[0-[2-[8-(phenylmethylamino)octylamino]ethyl]-oxime];

Erythromycin A (E)-9-[0-[2-[5-(phenylmethylamino)pentylamino]ethyl]-oxime];

Erythromycin A (E)-9-[0-[5-[6-(phenylmethylamino)hexylamino]pentyl]-oxime];

Erythromycin A (E)-9-[0-[3-[6-(phenylmethylamino)hexylamino]propyl]-oxime];

Erythromycin A (E)-9-[0-[3-[4-(phenylmethylamino)butylamino]propyl]-

oxime];

Erythromycin A (E)-9-[0-[2-[N-methyl-6-(N'-methyl-N'-phenylmethyl-amino)hexylamino]ethyl]oxime];

5 Erythromycin A (E)-9-[0-[2-[6-[(biphenyl-4-yl)methylamino]hexylamino]ethyl]oxime];

Erythromycin A (E)-9-[0-[2-[6-[(3-phenoxyphenyl)methylamino]hexylamino]ethyl]oxime];

Erythromycin A (E)-9-[0-[2-[6-[(4-phenoxyphenyl)methylamino]hexylamino]ethyl]oxime];

10 Erythromycin A (E)-9-[0-[2-[6-(2-furylmethylamino)hexylamino]ethyl]oxime];

Erythromycin A (E)-9-[0-[2-[6-(3-pyridylmethylamino)hexylamino]ethyl]oxime];

15 Erythromycin A (E)-9-[0-[2-[6-[(4-methoxyphenyl)methylamino]hexylamino]ethyl]oxime];

Erythromycin A (E)-9-[0-[2-[6-[(4-n.butoxyphenyl)methylamino]hexylamino]ethyl]oxime];

Erythromycin A (E)-9-[0-[2-[6-[(3,4-methylenedioxyphenyl)methylamino]hexylamino]ethyl]oxime];

20 Erythromycin A (E)-9-[0-[2-[6-[(4-methylsulphonylphenyl)methylamino]hexylamino]ethyl]oxime];

Erythromycin A (E)-9-[0-[2-[6-[(4-fluorophenyl)methylamino]hexylamino]ethyl]oxime];

Erythromycin A (E)-9-[0-[2-[6-[(2-trifluoromethylphenyl)methylamino]hexylamino]ethyl]oxime];

Erythromycin A (E)-9-[0-[2-[6-[(3-trifluoromethylphenyl)methylamino]hexylamino]ethyl]oxime];

Erythromycin A (E)-9-[0-[2-[6-[(4-trifluoromethylphenyl)methylamino]hexylamino]ethyl]oxime];

- 9 -

Erythromycin A (E)-9-[0-[2-[6-[(2-hydroxyphenyl)methylamino]hexyl-
amino]ethyl]oxime];

Erythromycin A (E)-9-[0-[2-[6-[(3-hydroxyphenyl)methylamino]hexyl-
5 amino]ethyl]oxime];

Erythromycin A (E)-9-[0-[2-[6-[(4-hydroxyphenyl)methylamino]hexyl-
amino]ethyl]oxime];

Erythromycin A (E)-9-[0-[2-[6-[(3,5-dichloro-2-hydroxyphenyl)methyl-
amino]hexylamino]ethyl]oxime];

Erythromycin A (E)-9-[0-[2-[6-[(2-nitrophenyl)methylamino]hexylami-
10 no]ethyl]oxime];

Erythromycin A (E)-9-[0-[2-[6-[(3-nitrophenyl)methylamino]hexylami-
no]ethyl]oxime];

Erythromycin A (E)-9-[0-[2-[6-[(4-nitrophenyl)methylamino]hexylami-
15 no]ethyl]oxime];

Erythromycin A (E)-9-[0-[2-[6-[(4-hydroxy-3-nitrophenyl)methylami-
no]hexylamino]ethyl]oxime];

Erythromycin A (E)-9-[0-[2-[6-[(3-hydroxy-4-nitrophenyl)methylami-
no]hexylamino]ethyl]oxime];

Erythromycin A (E)-9-[0-[2-[N-methyl-6-[N'-methyl-N'-(4-trifluoro-
20 methylphenyl)methylamino]hexylamino]ethyl]oxime].

The preparation of the compounds of formula (I), object of the
present invention, can be carried out according to the synthetic
method which is described below.

The method comprises, at first, the condensation reaction between a
25 suitable amino acid of formula



wherein

R_1 and m have the above reported meanings;


with an acyl chloride of formula



wherein

5 A and n have the above reported meanings.

The condensation reaction is carried out, according to conventional techniques, in an inert solvent and in the presence of a base such as, for instance, an alkali metal hydroxide, to obtain the compounds of formula



wherein

A, R₁, n and m have the above reported meanings.

The thus obtained N-acyl amino acids of formula (IV) are further condensed, according to conventional techniques, with an amino ester
15 of formula




wherein

20 R₂ and r have the above reported meanings;

R₄ represents a methyl or ethyl group;

to obtain the compounds of formula




wherein

25 A, R₁, R₂, R₄, n, m and r have the above reported meanings.

By working according to conventional techniques the compounds of formula (VI) are subsequently reduced, for instance with sodium boron hydride in the presence of acids, lithium aluminum hydride, dimethyl sulphide-borane or by catalytic hydrogenation, to the


corresponding amino alcohols of formula



5 wherein

A, R₁, R₂, n, m and r have the above reported meanings.

The amino alcohols of formula (VII) are then converted into the corresponding sulphonyl derivatives of formula (VIII), for instance by means of methanesulphonyl chloride or p.toluenesulphonyl chloride, and subsequently condensed with Erythromycin A 9-O-oxime or 6-O-methylerythromycin A 9-O-oxime, both representable according to formula (IX), to obtain the compounds of formula (I)



wherein

A, R₁, R₂, M, n, m and r have the above reported meanings;

20 R_s represents a mesyl or tosyl group.

The reaction between the compounds of formula (VIII) and the oximes of formula (IX) is carried out in an inert organic solvent such as, for instance, tetrahydrofuran, ethyl ether or 1,2-dimethoxyethane, in the presence of potassium tert-butyrate and of 18-crown-6 ether as complexing agent.

It is clear to the man skilled in the art that when the sulphonylation reaction is carried out by using the compounds of formula (VII) wherein one or both R₁ and R₂ substituents represent a hydrogen

atom, it can be necessary to protect the nitrogen atom or atoms, before carrying out the sulphonylation reaction.


In that case, the condensation of the thus obtained N-protected
 5 sulphonyl derivatives with the oximes of formula (IX), analogously to what previously reported, and the subsequent deprotection carried out according to conventional methods, allow to obtain the compounds of formula (I) wherein one or both R₁ and R₂ substituents represent a hydrogen atom.

10 For a bibliographic reference to the protection of amines see [T. W. Greene and P.G.M. Wuts, Protective groups in organic synthesis, John Wiley & Sons, Inc., 2nd. Ed., (1991), 309-405].

The compounds of formula (II), (III) and (V) are known or easily prepared according to known methods.

15 Also the oximes of formula (IX) are known compounds and can be prepared according to conventional methods comprising, for instance, the reaction of Erythromycin A or 6-O-methylerythromycin A with hydroxylamine hydrochloride.

The esters of formula (VI) can be optionally prepared according to
 20 an alternative synthetic method comprising, at first, the condensation of a suitable amino acid of formula (II) with an amino ester of formula (V), to obtain the compounds of formula



wherein

25 R₁, R₂, R₄, m and r, have the above reported meanings.

It is clear to the man skilled in the art that before carrying out the condensation between the amino acid of formula (II) and the amino ester of formula (V) it can be necessary to suitably protect according to what already reported for the sulphonylation reaction.


the amino group of the amino acid of formula (II).

The further condensation of the compounds of formula (X) with a compound of formula (III), carried out according to conventional techniques and the optional deprotection allow then to obtain the compounds of formula (VI).

The preparation of the compounds of formula (I) wherein at least one of the two R_1 and R_2 substituents represents a group selected among ethyl, n.propyl, n.butyl and isobutyl, can be carried out according to an alternative synthetic method which is described hereinafter.

Said method comprises, at first, the acylation of the nitrogen atom or atoms of the amino alcohols of formula (VII) wherein one or both R_1 and R_2 substituents represent a hydrogen atom.

For instance, by using a compound of formula (VII) wherein both R_1 and R_2 substituents represent a hydrogen atom and by working according to conventional techniques in the presence of a suitable acyl chloride ($R'COCl$), it is possible to obtain the compounds of formula




wherein

A , n , m and r have the above reported meanings;

R' represents a straight or branched C_1 - C_8 alkyl group.

The reduction of the compounds of formula (XI), carried out according to conventional methods, allows to obtain the compounds of formula




wherein

A , n , m and r have the above reported meanings;

R_1 and R_2 represent ethyl, n.propyl, n.butyl or isobutyl groups;

which, converted into the corresponding sulphonyl derivatives and condensed with the oximes of formula (IX), analogously to what previously reported, allow to obtain the compounds of formula




wherein

A, M, n, m and p have the above reported meanings;

R₁ and R₂ represent ethyl, n.propyl, n.butyl or isobutyl groups.

10 An alternative synthetic process with respect to those previously reported for the preparation of the compounds of formula (I), object of the present invention, is described hereinafter.

Said process comprises, at first, the oxidation of a suitable N-protected amino alcohol such as, for instance, an N-benzyloxycarbonyl-amino alcohol of formula (XIII), in the presence of sodium
 15 hypochlorite and of the free radical 2,2,6,6-tetramethylpiperidinoxy (TEMPO), in an inert organic solvent, to obtain the compounds of formula (XIV)



wherein


m has the above reported meanings;

Z represents a protecting group.

25 Examples of inert organic solvents usable in the oxidation reaction are, for instance, methylene chloride, chloroform, carbon tetrachloride, 1,2-dichloroethane, ethyl acetate, benzene and toluene.

The amination of the thus obtained aldehyde in the presence of a suitable amino alcohol of formula (XV) and the reduction of the formed intermediate, for instance in the presence of sodium boron

hydride, allow to obtain the amino alcohols of formula (XVI)




5 (XV) (XVI)

wherein

Z, m and r have the above reported meanings.


The further protection of the compounds of formula (XVI) at the amino nitrogen and, in this order, the conversion into the corresponding sulphonyl derivatives, the condensation with the oximes of
10 formula (IX) and the deprotection at the nitrogen atoms, analogously to what previously reported, allow to obtain the compounds of formula



wherein

M, m and r have the above reported meanings.

The intermediate oximes of formula (XVII), condensed with a suitable aldehyde of formula (XVIII) and reduced, for instance by catalytic
20 hydrogenation, allow to obtain the compounds of formula (I)



wherein

A, M, n, m and r have the above reported meanings.

25 The compounds of formula (XIII), (XV) and (XVIII) are known or easily prepared according to known methods.

The compounds of formula (I) wherein one or both R₁ and R₂ substituents represent a hydrogen atom, prepared according to one of the previously described methods, can be optionally alkylated at the

nitrogen atom or atoms of the di-amino moiety according to conventional methods comprising, for instance, the condensation with a suitable aldehyde and the reduction of the obtained intermediate.

- 5 The compounds of formula (I) wherein R_1 and R_2 , the same or different, represent a straight or branched C_1 - C_4 alkyl group are thus obtained.

The preparation of the compounds of formula (I) with Z or E configuration is carried out according to one of the synthetic schemes
10 above described, by using the oxime of formula (IX) in the desired configuration [J. C. Gasc et al., The Journal of Antibiotics, 44, 313-330, (1991)].

The compounds of formula (I) are endowed with antibacterial activity against several Gram-positive and Gram-negative microorganisms and
15 are useful in clinical and veterinary therapy for the treatment of several infectious diseases such as, for instance, the infections of the central nervous system, of the upper and lower respiratory tract, of the gastrointestinal tract, of the urogenital tract, of the odontological tissue and of the external organs such as skin,
20 eye and ear.

Said compounds, furthermore, resulted to be active with respect to several Gram-positive microorganisms of clinical interest, resistant to Erythromycin A or, more in general, to macrolide antibiotics characterized by the presence of a 14 or 15 members macrolactone.

The antibacterial activity of the compounds of formula (I) against
25 Gram-positive microorganisms such as Streptococcus pyogenes, Streptococcus pneumoniae, Enterococcus faecalis and Staphylococcus aureus and Gram-negative microorganisms such as Escherichia coli and Klebsiella pneumoniae was evaluated by means of in vitro tests suitable to evaluate the minimum concentration of antibiotic

enabling the inhibition of the bacterial growth (HIC) (example 23).
Roxithromycin and Clarithromycin were used as reference compounds
[The Merck Index, XI Ed., No. 8253 and 2340, respectively].

- 5 The antibacterial activity of the compounds of formula (I) against
Gram-positive microorganisms resulted to be practically comparable
to that of Roxithromycin and Clarithromycin, macrolides both
characterized by a high antibacterial activity in vitro (table 1).
With respect to Gram-negative microorganisms and, particularly,
10 against enterobacteria such as Escherichia coli and Klebsiella
pneumoniae, the compounds of formula (I) resulted to be markedly
more active than both reference compounds (table 2).

- To this extent, it is interesting to point out that the compounds of
formula (I), object of the present invention, resulted to be more
15 active than Roxithromycin, described in the aforementioned European
patent application No. 0033255 and chosen as election compound with
respect to several other derivatives such as, for instance,
Erythromycin A 9-[0-[(2-dimethylamino)ethyl]oxime] [J. C. Gasc et
al., The Journal of Antibiotics 44, 313-330, (1991)].

- 20 Moreover, the compounds of formula (I) resulted to be active in vivo
(table 3).

The in vivo antibacterial activity of the compounds of formula (I),
expressed as average protecting dose PD₅₀ (mg/Kg), was evaluated by
means of experimental pulmonary infection induced in mouse by
Streptococcus pyogenes (example 23).

- 25 By considering the data of activity in vivo it is evident that the
compounds of formula (I) are characterized by a prolonged duration
of action and a long half-life of tissue elimination.

In fact, after intraperitoneal administration in mouse, the com-
pounds of formula (I) are rapidly and widely distributed in the

whole organism and the tissue levels result to be higher than the plasmatic ones.

This results particularly evident by considering the PD_{50} values for
5 the compounds of formula (I) administered 24 hours before or 1 hour after the infection.

Said values, in fact, result to be substantially unchanged after administration 24 hours before or 1 hour after the infection.

In the case of experimental pulmonary infection induced in mouse by
10 *Streptococcus pyogenes*, a pathogen traditionally responsible for respiratory diseases, effective concentrations of the compounds of formula (I) intraperitoneally administered, persist at pulmonary level 24-48 hours after the administration.

The reference compounds Roxithromycin and Clarithromycin administered 24 hours before the infection, instead, resulted to be inactive.
15

Therefore, the compounds of formula (I) result also to be endowed with a pulmonary selectivity and can be advantageously used in the treatment of the infections of the respiratory tract.

In addition to the aforementioned activity against bacterial
20 microorganisms the compounds of formula (I), object of the present invention, resulted to be active against eukaryotic pathogens. In particular, they resulted to be markedly active against protozoa such as *Plasmodium falciparum* which is the well-known malaria parasite.

25 The compounds of formula (I), therefore, can also be advantageously used in the treatment of malarian diseases.

Besides being characterized by a wide spectrum of antibiotic activity against Gram-positive and Gram-negative microorganisms and protozoa, by a good stability to the acids and by good

pharmacokinetic properties, the compounds of formula (I) present, in mouse, an acute toxicity comparable to that of Roxithromycin.

Therefore, being characterized by a high safety of use, they can be
5 advantageously employed in human and veterinary therapy.

The compounds of formula (I) will be preferably used in a suitable pharmaceutical form useful to oral, parenteral, suppository or topical administration.

Object of the present invention, therefore, are the pharmaceutical
10 compositions containing a therapeutically effective amount of one or more compounds of formula (I) in admixture with a pharmaceutically acceptable carrier.

Said pharmaceutical forms comprise tablets, capsules, syrups, injectable solutions ready to use or to be prepared when used by
15 dilution of a lyophilized, suppositories, solutions, creams, ointments and eye lotions.

For the veterinary use, in addition to the above compositions, it is possible to prepare solid or liquid concentrates to be diluted into the feed or drinking water.

According to the type of composition, besides a therapeutically
20 effective amount of one or more compounds of formula (I), they will contain solid or liquid excipients or diluents for pharmaceutical or veterinary use and optionally other additives of normal formulating use such as thickeners, aggregants, lubricants, disgregants, flavouring and colouring agents.

25 In order to treat particular infections, the compound of formula (I) could be in association with an effective amount of another active ingredient.

The effective amount of the compound of formula (I) can vary according to different factors such as the seriousness and the phase of

the infection, the organ or the system affected, the characteristics of the host species, the susceptibility of the bacterial species responsible for the infection and the selected route of administration.

The therapeutic dose will be generally comprised between 0.5 and 100 mg/Kg of body weight/day and could be administered into a single dose or into more daily doses.

With the aim of illustrating the present invention, without limiting it, the following examples are now given.

The structures of the compounds of formula (I) and of the synthetic intermediates for their preparation were confirmed by ^1H -NMR or by ^{13}C -NMR spectroscopy. The values of the meaningful signals of the more advanced intermediates and of the compounds of formula (I) are reported hereinafter.

15

Example 1

Preparation of N-benzoyl-6-aminohexanoic acid

A solution of benzoyl chloride (0.18 moles) in ethyl ether (160 ml) and a solution of sodium hydroxide 1N (180 ml) were contemporaneously added to a mixture of 6-aminohexanoic acid (0.15 moles) in ethyl ether (150 ml) and water (200 ml) containing sodium hydroxide (0.15 moles), kept under stirring at a temperature comprised between 0-5°C.

At the end of the additions, the reaction mixture was brought at room temperature and kept under stirring for other 4 hours.

After separation of the phases, the aqueous phase was washed with ethyl ether (200 ml) and acidified at Congo red with concentrated hydrochloric acid.

After extraction with ethyl acetate (3x200 ml) the collected organic phases were washed with a saturated aqueous solution of sodium

30

chloride (200 ml), dried on sodium sulphate and evaporated at reduced pressure.

5 N-benzoyl-6-aminohexanoic acid, used as such in the subsequent reactions, was thus obtained.

By working analogously the following compounds were prepared:

N-benzoyl-3-aminopropanoic acid;

N-benzoyl-glycine;

N-benzoyl-8-aminooctanoic acid;

10 N-benzoyl-4-aminobutanoic acid;

N-phenylacetyl-6-aminohexanoic acid;

N-phenylacetyl-glycine;

N-benzoyl-N-isopropyl-4-aminobutanoic acid;

N-benzoyl-N-isopropyl-6-aminohexanoic acid.

Example 2

15

Preparation of N-[6-(benzoylamino)hexanoyl]glycine ethyl ester

A solution of dicyclohexylcarbodiimide (112 mmoles) in anhydrous tetrahydrofuran (44 ml) was gradually added to a suspension of N-benzoyl-6-aminohexanoic acid (93.5 mmoles), prepared as described in example 1, glycine ethyl ester hydrochloride (112 mmoles), tri-
20 ethylamine (112 mmoles) and anhydrous 1-hydroxybenzotriazole (112 mmoles) in tetrahydrofuran (330 ml), kept under stirring at 0°C.

The reaction mixture was brought at room temperature and kept under stirring for 16 hours.

25 At the end, a precipitate was formed which was eliminated by filtration and the thus obtained filtrate was evaporated at reduced pressure.

The residue was collected with ethyl acetate (300 ml) and subsequently washed with a solution of hydrochloric acid at 5% (2x100 ml), with a saturated solution of sodium chloride (100 ml), with a

30

- 22 -

solution of sodium bicarbonate at 5% (2x100 ml) and, at last, with a saturated solution of sodium chloride (100 ml).

The organic phase was dried on sodium sulphate and evaporated to dryness at reduced pressure, thus obtaining N-[6-(benzoylamino)hexanoyl]glycine ethyl ester which was used as such in the subsequent reactions.

By working analogously the following compounds were prepared:

- N-[(benzoylamino)acetyl]glycine ethyl ester;
N-[6-(phenylacetylamino)hexanoyl]glycine ethyl ester;
N-[(phenylacetylamino)acetyl]glycine ethyl ester;
ethyl 6-[6-(benzoylamino)hexanoylamino]hexanoate;
N-[5-(benzoylamino)pentanoyl]glycine methyl ester;
methyl 6-[5-(benzoylamino)pentanoylamino]hexanoate;
N-[7-(benzoylamino)heptanoyl]glycine methyl ester;
methyl 5-[6-(benzoylamino)hexanoylamino]pentanoate;
methyl 6-[(benzoylamino)acetylaminol]hexanoate;
methyl 3-[6-(benzoylamino)hexanoylamino]propionate;
ethyl 6-[N-isopropyl-(phenylacetylamino)acetylaminol]hexanoate;
methyl 6-[4-(benzoylamino)butanoylamino]hexanoate;
methyl 4-[N-isopropyl-4-(N'-isopropyl-N'-benzoylamino)butanoylami-
nol]butanoate.

Example 3

Preparation of N-(6-aminohexanoyl)glycine ethyl ester

a) 6-Aminohexanoic acid (100 g; 0.762 moles) and, gradually, a solution of di-tert-butyl dicarbonate (168 g; 0.762 moles) in methanol (140 ml) were added to a solution of sodium hydroxide (33.54 g; 0.831 moles) in water (840 ml) and methanol (400 ml).

The reaction mixture was kept under stirring at room temperature for 4 hours.

After that, solid di-tert-butyl dicarbonate (17.5 g; 0.08 moles) was added again, keeping under stirring for other 16 hours.

The reaction mixture was then washed with hexane (2x400 ml), acidified up to pH 1.5 with a solution of potassium bisulphate and extracted with ethyl acetate (3x450 ml).

The collected organic phases were dried on sodium sulphate and evaporated to dryness affording thus 6-(tert-butoxycarbonylamino)-hexanoic acid as an oil (163 g).

b) By working analogously to what described in example 2, 6-(tert-butoxycarbonylamino)hexanoic acid (163 g) was directly condensed with glycine ethyl ester hydrochloride (118 g; 0.845 moles), obtaining thus N-[6-(tert-butoxycarbonylamino)hexanoyl]glycine ethyl ester (285 g) as a raw product which was used as such in the subsequent reaction.

m.p. 76-77°C (isopropyl ether)

c) A solution of hydrochloric acid 6N (150 ml) in ethyl acetate (150 ml) was added to a solution of N-[6-(tert-butoxycarbonylamino)hexanoyl]glycine ethyl ester (285 g) in ethyl acetate (500 ml), kept under stirring at room temperature.

After 24 hours a precipitate was formed which was filtered, washed with ethyl acetate and with ethyl ether, and dried in oven (50°C) under vacuum.

N-(6-aminohexanoyl)glycine ethyl ester (93 g) was thus obtained as a raw product which was used as such in the subsequent reactions.

TLC (methylene chloride:methanol:ammonia=10:2:1) Rf=0.2.

Example 4

Preparation of N-[6-[(4-fluorobenzoyl)aminohexanoyl]glycine ethyl ester

A solution of 4-fluorobenzoyl chloride (47.4 mmoles) in methylene

chloride (30 ml) was gradually added to a suspension of N-(6-amino-hexanoyl)glycine ethyl ester (39.5 mmoles), prepared as described in example 3, and triethylamine (87 mmoles) in methylene chloride (150 ml), kept under stirring at 0°C.

The thus prepared mixture, to which triethylamine (2 ml) was subsequently added, was brought at room temperature and kept under stirring.

After one hour under these conditions, the reaction mixture was washed with a solution of hydrochloric acid at 5% (2x100 ml) and with a saturated solution of sodium chloride (3x100 ml).

The separated organic phase was dried on sodium sulphate and evaporated to dryness under vacuum.

N-[6-[(4-fluorobenzoyl)amino]hexanoyl]glycine ethyl ester was thus obtained as a crude product, used as such in the subsequent reactions.

m.p. 121-122°C (ethyl acetate)

TLC (ethyl acetate) Rf=0.3

By working analogously the following compound was prepared:

N-[6-(2-furoylamino)hexanoyl]glycine ethyl ester

m.p. 104-106°C (acetonitrile/isopropyl ether)

TLC (methylene chloride:methanol=95:5) Rf=0.3.

Example 5

Preparation of N-[6-[(4-methoxybenzoyl)amino]hexanoyl]glycine ethyl ester

By working analogously to what described in example 2 and by using 4-methoxybenzoic acid (33 mmoles) and N-(6-aminohexanoyl)glycine ethyl ester (39.5 mmoles), prepared as described in example 3, N-[6-[(4-methoxybenzoyl)amino]hexanoyl]glycine ethyl ester was obtained as a crude product, used as such in the subsequent

reactions.

m.p. 106-107°C

TLC (methylene chloride:methanol=90:10) Rf=0.46

5 By working analogously the following compounds were prepared:

N-[6-[(3,4-methylenedioxybenzoyl)amino]hexanoyl]glycine ethyl ester

TLC (methylene chloride:methanol=90:10) Rf=0.39;

N-[6-[(4-methylsulphonylbenzoyl)amino]hexanoyl]glycine ethyl ester

m.p. 124-126°C

TLC (methylene chloride:methanol=96:4) Rf=0.31;

10 N-[6-[(3-trifluoromethylbenzoyl)amino]hexanoyl]glycine ethyl ester

m.p. 102-104°C

TLC (methylene chloride:methanol=95:5) Rf=0.38.

Example 6

15 Preparation of 2-[6-(phenylmethylamino)hexylamino]ethanol

Sulphuric acid 6N in ethyl ether (40.9 ml; 700 mmoles), prepared by mixing sulphuric acid 96% (33 ml) and ethyl ether (100 ml), was gradually added to a suspension of N-[6-(benzoylamino)hexanoyl]glycine ethyl ester (46.8 mmoles), prepared as described in example 2, and sodium boron hydride (700 mmoles) in anhydrous tetrahydrofuran (200 ml), kept under stirring at a temperature comprised between 15°C and 20°C.

The reaction mixture was brought at boiling temperature for 24 hours and, subsequently, cooled at 0°C.

Methanol (150 ml) was then added under stirring.

25 The solvent was evaporated at reduced pressure and the residue was collected with a solution of sodium hydroxide 6N (200 ml), keeping the resultant mixture at the boiling temperature for 24 hours.

The reaction mixture, cooled at room temperature, was then extracted with tetrahydrofuran (2x100 ml) and the organic phase was evaporated

to dryness, collected with ethyl acetate and dried on sodium sulphate.

Through acidification with an ethereal solution of hydrochloric acid
5 a precipitate constituted by 2-[6-(phenylmethvlamino)hexylamino]-
ethanol as hydrochloride salt was obtained.

The thus obtained product was used as such in the subsequent reactions.

By working analogously the following compounds were prepared:

- 10 2-[2-(phenylmethvlamino)ethylamino]ethanol;
- 2-[6-(2-phenylethvlamino)hexylamino]ethanol;
- 6-[6-(phenylmethvlamino)hexylamino]hexanol;
- 2-[5-(phenylmethvlamino)pentylamino]ethanol;
- 2-[8-(phenylmethvlamino)octylamino]ethanol;
- 15 5-[6-(phenylmethvlamino)hexylamino]pentanol;
- 6-[3-(phenylmethvlamino)propylamino]hexanol;
- 3-[6-(phenylmethvlamino)hexylamino]propanol;
- 3-[4-(phenylmethvlamino)butylamino]propanol;
- 6-[2-(phenylmethvlamino)ethylamino]hexanol;
- 20 6-[N-isopropyl-4-(phenylmethvlamino)butylamino]hexanol;
- 2-[6-[(4-fluorophenyl)methylamino]hexylamino]ethanol;
- 2-[6-[(4-methoxyphenyl)methylamino]hexylamino]ethanol;
- 2-[6-[(3,4-methylenedioxyphenyl)methylamino]hexylamino]ethanol;
- 2-[6-[(3-trifluoromethylphenyl)methylamino]hexylamino]ethanol;
- 2-[6-[(4-methylsulphonylphenyl)methylamino]hexylamino]ethanol;
- 25 4-[N-isopropyl-4-(N'-isopropyl-N'-phenylmethvlamino)butylamino]butanol.

Example 7

Preparation of 6-[N-acetyl-6-(N'-acetyl-N'-phenylmethvlamino)hexyl-
aminol]hexyl acetate

Triethylamine (1.95 ml; 14 mmoles) and a solution of acetyl chloride (0.62 ml; 8.69 mmoles) in methylene chloride (5 ml) were gradually added to a suspension of 6-[6-(phenylmethylamino)hexylamino]hexanol
5 (1 g; 2.6 mmoles), prepared as described in example 6, in methylene chloride (15 ml), kept under stirring at 0°C.

After one hour under stirring at 0°C, the reaction mixture was brought at room temperature and kept under stirring for other 16 hours.

10 The reaction mixture was then washed with a solution of hydrochloric acid at 10% (10 ml) and with a saturated solution of sodium chloride.

After separation of the phases, the organic phase was dried on sodium sulphate and evaporated to dryness under vacuum, obtaining
15 thus 6-[N-acetyl-6-(N'-acetyl-N'-phenylmethylamino)hexylamino]hexyl acetate (1.18 g) as an oil, used as such in the subsequent reactions.

By working analogously the following compound was prepared:

2-[N-acetyl-6-[N'-acetyl-N'-(2-phenylethyl)aminohexylamino]ethyl
20 acetate.

Example 8

Preparation of 6-[N-ethyl-6-(N'-ethyl-N'-phenylmethylamino)hexylamino]hexanol

By working analogously to what described in example 6 and by using
25 6-[N-acetyl-6-(N'-acetyl-N'-phenylmethylamino)hexylamino]hexyl acetate, prepared as described in example 7, 6-[N-ethyl-6-(N'-ethyl-N'-phenylmethylamino)hexylamino]hexanol was prepared.

By working analogously the following compound was prepared:

2-[N-ethyl-6-[N'-ethyl-N'-(2-phenylethyl)aminohexylamino]ethanol.

Example 9

Preparation of 2-[N-benzyloxycarbonyl-6-(N'-benzyloxycarbonyl-N'-phenylmethylamino)hexylamino]ethanol

A solution of sodium hydroxide 1N (44.5 ml) and a solution in
5 toluene at 50% of benzyl chloroformate (44.5 mmoles) in ethyl
acetate (33 ml) were gradually and contemporaneously added to a
solution of 2-[6-(phenylmethylamino)hexylamino]ethanol dihydrochloride (18.5 mmoles), prepared as described in example 6, in a solution of sodium hydroxide 1N (37.1 ml) and ethyl acetate (40 ml),
10 kept under stirring at a temperature of 0°C.

At the end of the additions, the reaction mixture was brought at
room temperature and kept under stirring for 24 hours.

After separation of the phases, the aqueous phase was washed with
ethyl acetate (2x50 ml).

15 The collected organic phases were washed with a saturated solution
of sodium chloride (50 ml), dried on sodium sulphate and evaporated
to dryness under vacuum.

2-[N-benzyloxycarbonyl-6-(N'-benzyloxycarbonyl-N'-phenylmethylamino)hexylamino]ethanol was thus obtained as an oil, used as such in
the subsequent reactions.

20 TLC (ethyl acetate:hexane=50:50) Rf=0.20.

By working analogously the following compounds were prepared:

2-[N-benzyloxycarbonyl-2-(N'-benzyloxycarbonyl-N'-phenylmethylamino)ethylamino]ethanol

TLC (ethyl acetate:hexane=60:40) Rf=0.25;

25 6-[N-benzyloxycarbonyl-6-(N'-benzyloxycarbonyl-N'-phenylmethylamino)hexylamino]hexanol

TLC (ethyl acetate:hexane=50:50) Rf=0.27;

6-[N-benzyloxycarbonyl-5-(N'-benzyloxycarbonyl-N'-phenylmethylamino)pentylamino]hexanol;

- 2-[N-benzoyloxycarbonyl-5-(N'-benzoyloxycarbonyl-N'-phenylmethy-
lamino)pentylamino]ethanol;
- 2-[N-benzoyloxycarbonyl-8-(N'-benzoyloxycarbonyl-N'-phenylmethy-
lamoctylamino]ethanol;
- 5 5-[N-benzoyloxycarbonyl-6-(N'-benzoyloxycarbonyl-N'-phenylmethy-
lamino)hexylamino]pentanol;
- 6-[N-benzoyloxycarbonyl-3-(N'-benzoyloxycarbonyl-N'-phenylmethy-
lamino)propylamino]hexanol;
- 10 3-[N-benzoyloxycarbonyl-6-(N'-benzoyloxycarbonyl-N'-phenylmethy-
lamino)hexylamino]propanol;
- 3-[N-benzoyloxycarbonyl-4-(N'-benzoyloxycarbonyl-N'-phenylmethy-
lamino)butylamino]propanol;
- 6-[N-isopropyl-2-[N'-benzoyloxycarbonyl-N'-(2-phenylethyl)amino]eth-
ylamino]hexanol
- 15 TLC (methylene chloride:methanol:ammonia=95:5:0.5) Rf=0.33;
- 6-[N-benzoyloxycarbonyl-4-(N'-isopropyl-N'-phenylmethy-
lamino)butylamino]hexanol
- TLC (methylene chloride:methanol:ammonia=95:5:0.5) Rf=0.42;
- 20 2-[N-benzoyloxycarbonyl-6-[N'-benzoyloxycarbonyl-N'-[(4-fluorophenyl)-
methyl]amino]hexylamino]ethanol
- TLC (ethyl acetate:hexane=60:40) Rf=0.35;
- 2-[N-benzoyloxycarbonyl-6-[N'-benzoyloxycarbonyl-N'-[(4-methoxyphen-
yl)methyl]amino]hexylamino]ethanol
- TLC (ethyl acetate:hexane=50:50) Rf=0.2;
- 25 2-[N-benzoyloxycarbonyl-6-[N'-benzoyloxycarbonyl-N'-[(3,4-methylenedi-
oxyphenyl)methyl]amino]hexylamino]ethanol
- TLC (ethyl acetate:hexane=60:40) Rf=0.26;
- 2-[N-benzoyloxycarbonyl-6-[N'-benzoyloxycarbonyl-N'-[(3-trifluorometh-
ylphenyl)methyl]amino]hexylamino]ethanol

TLC (ethyl acetate:hexane=50:50) Rf=0.25;

2-[N-benzyloxycarbonyl-6-(N'-benzyloxycarbonyl-N'-(4-methylsulphonylphenyl)methylamino)hexylamino]ethanol

5 TLC (ethyl acetate:hexane=90:10) Rf=0.36.

Example 10

Preparation of 2-[N-benzyloxycarbonyl-6-(N'-benzyloxycarbonyl-N'-phenylmethylamino)hexylamino]ethyl-methanesulphonate

A solution of methanesulphonyl chloride (3.16 mmoles) in methylene chloride (5 ml) was gradually added to a solution of 2-[N-benzyloxy-
10 carbonyl-6-(N'-benzyloxycarbonyl-N'-phenylmethylamino)hexylamino]-ethanol (2.6 mmoles), prepared as described in example 9, in methylene chloride (15 ml) containing triethylamine (0.44 ml; 3.16 mmoles), under stirring and at a temperature of 0°C.

15 The reaction mixture, brought at room temperature and kept under stirring for 5 hours, was added to a solution of hydrochloric acid at 5% (20 ml).

After separation of the phases, the organic phase was washed with hydrochloric acid at 5% (10 ml) and with a saturated solution of
20 sodium chloride (3x10 ml).

The organic phase was then dried on sodium sulphate and evaporated to dryness affording thus 2-[N-benzyloxycarbonyl-6-(N'-benzyloxycarbonyl-N'-phenylmethylamino)hexylamino]ethyl-methanesulphonate, used as such in the reaction of the following example.

Example 11

25 Erythromycin A (E)-9-[0-[2-[N-benzyloxycarbonyl-6-(N'-benzyloxycarbonyl-N'-phenylmethylamino)hexylamino]ethyl]oxime]

Erythromycin A (E)-9-O-oxime (627 mg; 0.84 mmoles), 18-crown-6 ether (220 mg; 0.84 mmoles) and a solution of 2-[N-benzyloxycarbonyl-6-(N'-benzyloxycarbonyl-N'-phenylmethylamino)hexylamino]ethyl-meth-

anesulphonate (0.84 mmoles), prepared as described in example 10, in anhydrous tetrahydrofuran (5 ml) were respectively added to a suspension of potassium tert-butyrate (103 mg; 0.92 mmoles) in anhydrous tetrahydrofuran (5 ml), kept at room temperature under stirring and nitrogen atmosphere.

The reaction mixture was kept under stirring at room temperature for 20 hours and, subsequently, evaporated at reduced pressure.

The residue was collected with ethyl acetate (10 ml) and the thus obtained mixture was washed with a saturated solution of sodium chloride (10 ml).

The aqueous phase was extracted with ethyl acetate (2x10 ml) and the collected organic phases were dried on sodium sulphate and evaporated to dryness.

15 Erythromycin A (E)-9-[0-[2-[N-benzyloxycarbonyl-6-(N'-benzyloxycarbonyl-N'-phenylmethylamino)hexylamino]ethyl]oxime] was thus obtained and used as such in the subsequent reactions.

TLC (methylene chloride:methanol:ammonia=90:9:1) Rf=0.58

Mass (C.I.) (M+H)⁺=1250;

20 ¹H-NMR (200 MHz, CDCl₃): δ (ppm): 7.38-7.10 (m, 15H, aromatics); 5.18-5.10 (m, 4H, 2 CH₂Ph); 3.30 (s, 3H, OCH₃); 2.26 (s, 6H, 2 NCH₃); 0.81 (t, 3H, CH₂CH₃).

By working analogously the following compounds were prepared:

25 Erythromycin A (E)-9-[0-[2-[N-benzyloxycarbonyl-2-(N'-benzyloxycarbonyl-N'-phenylmethylamino)ethylamino]ethyl]oxime]

TLC (methylene chloride:methanol:ammonia=90:10:1) Rf=0.5;

¹H-NMR (200 MHz, CDCl₃): δ (ppm): 7.11-6.97 (m, 15H, aromatics); 5.18-4.97 (m, 4H, 2 CH₂Ph); 3.30 (s, 3H, OCH₃); 2.25 (s, 6H, 2 NCH₃); 0.82 (t, 3H, CH₂CH₃);

30 Erythromycin A (E)-9-[0-[6-[N-benzyloxycarbonyl-6-(N'-benzyloxycar-

bonyl-N'-phenylmethylamino)hexylaminohexylloximel

TLC (methylene chloride:methanol:ammonia:90:10:1) Rf=0.6;

¹H-NMR (200 MHz, CDCl₃): δ (ppm): 7.27-6.96 (m, 15H, aromatics);

5 5.05-4.92 (m, 4H, 2 CH₂Ph); 3.17 (s, 3H, OCH₃); 2.13 (s, 6H, 2 NCH₃); 0.70 (t, 3H, CH₃CH₂);

Erythromycin A (E)-9-[0-[6-[N-benzyloxycarbonyl-3-(N'-benzyloxycarbonyl-N'-phenylmethylamino)propylaminohexylloximel

TLC (methylene chloride:methanol:ammonia:90:10:1) Rf=0.65;

10 Mass (C.I.) (M+H)⁺=1194;

¹H-NMR (200 MHz, CDCl₃): δ (ppm): 7.39-7.01 (m, 15H, aromatics);

5.17-5.02 (m, 4H, 2 CH₂Ph); 3.30 (s, 3H, OCH₃); 2.27 (s, 6H, 2 NCH₃); 0.82 (t, 3H, CH₃CH₂);

Erythromycin A (E)-9-[0-[6-(N-benzyloxycarbonyl-5-(N'-benzyloxycarbonyl-N'-phenylmethylamino)pentylaminohexylloximel;

15 bonyl-N'-phenylmethylamino)pentylaminohexylloximel;

Erythromycin A (E)-9-[0-[2-[N-benzyloxycarbonyl-8-(N'-benzyloxycarbonyl-N'-phenylmethylamino)octylaminoethylloximel;

bonyl-N'-phenylmethylamino)octylaminoethylloximel;

Erythromycin A (E)-9-[0-[2-[N-benzyloxycarbonyl-5-(N'-benzyloxycarbonyl-N'-phenylmethylamino)pentylaminoethylloximel;

bonyl-N'-phenylmethylamino)pentylaminoethylloximel;

20 Erythromycin A (E)-9-[0-[5-[N-benzyloxycarbonyl-6-(N'-benzyloxycarbonyl-N'-phenylmethylamino)hexylaminopentylloximel;

bonyl-N'-phenylmethylamino)hexylaminopentylloximel;

Erythromycin A (E)-9-[0-[3-[N-benzyloxycarbonyl-6-(N'-benzyloxycarbonyl-N'-phenylmethylamino)hexylaminopropylloximel;

bonyl-N'-phenylmethylamino)hexylaminopropylloximel;

Erythromycin A (E)-9-[0-[3-[N-benzyloxycarbonyl-4-(N'-benzyloxycarbonyl-N'-phenylmethylamino)butylaminopropylloximel;

25 bonyl-N'-phenylmethylamino)butylaminopropylloximel;

Erythromycin A (E)-9-[0-[6-[N-benzyloxycarbonyl-2-[N'-benzyloxycarbonyl-N'-(2-phenylethyl)aminoethylaminohexylloximel

bonyl-N'-(2-phenylethyl)aminoethylaminohexylloximel

m.p. 74-76°C

Mass (C.I.) (M+H)⁺=1172

¹H-NMR (200 MHz, CDCl₃): δ (ppm): 7.38-7.03 (m, 10H, aromatics);

5.13-5.03 (m, 2H, CH₂Ph); 3.29 (s, 3H, OCH₃); 2.25 (s, 6H, 2 NCH₃);

Erythromycin A (E)-9-[0-[6-[N-ethyl-6-(N'-ethyl-N'-phenylmethylamino)hexylamino]hexyl]oxime] (Compound 1)

5 m.p. 80-82°C (acetonitrile)

Mass (C.I.) (M+H)⁺=1094

¹³C-NMR (50 MHz, CDCl₃): δ (ppm): 175.20; 171.35; 140.06; 128.86; 128.07; 126.62; 102.96; 96.27; 53.54;

Erythromycin A (E)-9-[0-[2-[N-ethyl-6-[N'-ethyl-N'-(2-phenylethyl)-

10 aminohexylamino]ethyl]oxime] (Compound 2)

TLC (chloroform:hexane:triethylamine=45:45:10) R_f=0.2

Mass (C.I.) (M+H)⁺=1052

¹H-NMR (200 MHz, CDCl₃): δ (ppm): 7.26-7.04 (m, 5H, aromatics); 3.22 (s, 3H, OCH₃); 2.20 (s, 6H, 2 NCH₃); 0.79 (t, 3H, CH₂CH₃);

15 Erythromycin A (E)-9-[0-[6-[N-benzyloxycarbonyl-4-(N'-isopropyl-N'-phenylmethylamino)butylamino]hexyl]oxime]

m.p. 75-77°C

¹H-NMR (200 MHz, CDCl₃): δ (ppm): 7.47-7.12 (m, 10H, aromatics); 5.18-4.97 (m, 4H, 2 CH₂Ph); 3.30 (s, 3H, OCH₃); 2.25 (s, 6H, 2 NCH₃); 0.82 (t, 3H, CH₂CH₃);

20 Erythromycin A (E)-9-[0-[2-[N-benzyloxycarbonyl-6-[N'-benzyloxycarbonyl-N'-[(4-fluorophenyl)methyl]amino]hexylamino]ethyl]oxime]

TLC (methylene chloride:methanol:ammonia=90:10:1) R_f=0.62

25 ¹H-NMR (200 MHz, CDCl₃): δ (ppm): 7.38-6.88 (m, 15H, aromatics); 5.17-5.03 (m, 2H, CH₂Ph); 3.29 (s, 3H, OCH₃); 2.26 (s, 6H, 2 NCH₃); 0.81 (t, 3H, CH₂CH₃);

Erythromycin A (E)-9-[0-[2-[N-benzyloxycarbonyl-6-[N'-benzyloxycarbonyl-N'-[(4-methoxyphenyl)methyl]amino]hexylamino]ethyl]oxime]

TLC (methylene chloride:methanol:ammonia=45:45:10) R_f=0.3

30 ¹H-NMR (200 MHz, CDCl₃): δ (ppm): 7.40-7.23 (m, 10H, 2 PhCH₂O);

- 34 -

7.20-6.75 (m, 4H, PhOCH_3); 5.52-5.17 (m, 4H, 2 CH_2Ph); 3.77 (s, 3H, PhOCH_3); 3.29 (s, 3H, OCH_3); 2.25 (s, 6H, 2 NCH_3); 0.82 (t, 3H, CH_2CH_3);

- 5 Erythromycin A (E)-9-[0-[2-[N-benzoyloxycarbonyl-6-[N'-benzyloxycarbonyl-N'-[(3,4-methylenedioxyphenyl)methyl]amino]hexylamino]ethyl]-oximel

TLC (methylene chloride:methanol:ammonia=95:5:0.5) $R_f=0.31$

- $^1\text{H-NMR}$ (200 MHz, CDCl_3): δ (ppm): 7.38-7.22 (m, 10H, 2 PhCH_2O);
 10 6.78-6.55 (m, 3H, aromatics); 5.90 (s, 2H, OCH_2O); 5.15-5.02 (m, 4H, 2 CH_2Ph); 3.29 (s, 3H, OCH_3); 2.26 (s, 6H, 2 NCH_3); 0.82 (t, 3H, CH_2CH_3);

Erythromycin A (E)-9-[0-[2-[N-benzoyloxycarbonyl-6-[N'-benzyloxycarbonyl-N'-[(3-trifluoromethylphenyl)methyl]amino]hexylamino]ethyl]-

- 15 oximel

TLC (methylene chloride:methanol:ammonia=90:10:1) $R_f=0.65$

$^1\text{H-NMR}$ (200 MHz, CDCl_3): δ (ppm): 7.54-7.15 (m, 14H, aromatics);
 5.20-5.03 (m, 4H, 2 CH_2Ph); 3.30 (s, 3H, OCH_3); 2.26 (s, 6H, 2 NCH_3); 0.82 (t, 3H, CH_2CH_3);

- 20 Erythromycin A (E)-9-[0-[2-[N-benzoyloxycarbonyl-6-[N'-benzyloxycarbonyl-N'-[(4-methylsulphonylphenyl)methyl]amino]hexylamino]ethyl]-oximel

TLC (methylene chloride:methanol:ammonia=95:5:0.5) $R_f=0.5$

- $^1\text{H-NMR}$ (200 MHz, CDCl_3): δ (ppm): 7.90-7.79 (m, 4H, PhSO_2CH_3);
 25 7.48-7.15 (m, 10H, 2 PhCH_2O); 5.19-5.03 (m, 4H, 2 CH_2Ph); 3.30 (s, 3H, OCH_3); 3.02 (s, 3H, CH_3SO_2); 2.27 (s, 6H, 2 NCH_3); 0.82 (t, 3H, CH_2CH_3);

Erythromycin A (E)-9-[0-[4-[N-isopropyl-4-(N'-isopropyl-N'-phenylmethylamino)butylamino]butyl]oximel (Compound 3)

m.p. 83-85°C (hexane)

- 35 -

Mass (C.I.) (M+H)⁺=1066

¹H-NMR (200 MHz, CDCl₃): δ (ppm): 7.37-7.10 (m, 5H, aromatics); 3.50 (s, 2H, CH₂Ph); 3.30 (s, 3H, OCH₃); 2.26 (s, 6H, 2 NCH₃); 0.82 (t, 5 3H, CH₂CH₃).

Example 12

Preparation of Erythromycin A (E)-9-[0-[2-[6-(phenylmethylamino)hexylamino]ethyl]oxime] (Compound 4)

Palladium on charcoal at 10% (750 mg) was added to a solution of
10 Erythromycin A (E)-9-[0-[2-[N-benzyloxycarbonyl-6-(N'-benzyloxycarbonyl-N'-phenylmethylamino)hexylamino]ethyl]oxime] (5.9 mmoles), prepared as described in example 11, in ethanol (150 ml).

The thus prepared mixture was placed into a Parr hydrogenator loaded with hydrogen (1 bar) and kept under stirring at room temperature.

15 After 7 hours the catalyst was filtered off and the alcoholic solution was evaporated to dryness.

Erythromycin A (E)-9-[0-[2-[6-(phenylmethylamino)hexylamino]ethyl]oxime], purified by silica gel chromatography (eluent methylene chloride:methanol:ammonia=90:10:1) was thus obtained.

20 Mass (C.I.) (M+H)⁺=982

¹³C-NMR (50 MHz, CDCl₃): δ (ppm): 140.48; 128.39; 128.11; 126.88.

By working analogously the following compounds were prepared:

Erythromycin A (E)-9-[0-[2-[2-(phenylmethylamino)ethylamino]ethyl]oxime] (Compound 5)

25 ¹³C-NMR (50 MHz, CDCl₃): δ (ppm): 176.51; 172.36; 140.96; 129.08; 128.95; 127.67; 103.84; 96.86; 53.35;

Erythromycin A (E)-9-[0-[6-[6-(phenylmethylamino)hexylamino]hexyl]oxime] (Compound 6)

Mass (C.I.) (M+H)⁺=1038

30 ¹³C-NMR (50 MHz, CDCl₃): δ (ppm): 175.24; 171.31; 140.33; 128.37;

128.13; 126.89; 102.92; 96.27; 54.01;

Erythromycin A (E)-9-[0-[6-[3-(phenylmethylamino)propylamino]hexyl]-oxime] (Compound 7)

5 Mass (C.I.) (M+H)⁺=995

¹³C-NMR (50 MHz, CDCl₃): δ (ppm): 175.15; 171.37; 140.41; 128.38; 128.09; 126.89; 102.92; 96.27; 54.04;

Erythromycin A (E)-9-[0-[6-[5-(phenylmethylamino)pentylamino]hexyl]-oxime] (Compound 8)

10 ¹H-NMR (200 MHz, CDCl₃): δ (ppm): 7.35-7.15 (m, 5H, aromatics); 3.75 (s, 2H, CH₂Ph); 2.25 (s, 6H, 2 NCH₃); 0.81 (t, 3H, CH₂CH₃);

Erythromycin A (E)-9-[0-[2-[8-(phenylmethylamino)octylamino]ethyl]-oxime] (Compound 9)

15 ¹H-NMR (200 MHz, CDCl₃): δ (ppm): 7.40-7.15 (m, 5H, aromatics); 3.75 (s, 2H, CH₂Ph); 3.29 (s, 3H, OCH₃); 2.25 (s, 6H, 2 NCH₃); 0.82 (t, 3H, CH₂CH₃);

Erythromycin A (E)-9-[0-[2-[5-(phenylmethylamino)pentylamino]ethyl]-oxime] (Compound 10);

20 ¹³C-NMR (50 MHz, CDCl₃): δ (ppm): 174.96; 172.00; 140.31; 128.39; 128.14; 126.93; 103.16; 96.20; 53.98;

Erythromycin A (E)-9-[0-[5-[6-(phenylmethylamino)hexylamino]pentyl]-oxime] (Compound 11);

¹³C-NMR (50 MHz, CDCl₃): δ (ppm): 175.24; 171.24; 140.41; 128.38; 128.13; 126.88; 102.97; 96.28; 54.06;

25 Erythromycin A (E)-9-[0-[3-[6-(phenylmethylamino)hexylamino]propyl]-oxime] (Compound 12);

¹³C-NMR (50 MHz, CDCl₃): δ (ppm): 175.23; 171.46; 140.45; 128.39; 128.13; 126.88; 102.99; 96.29; 50.06;

Erythromycin A (E)-9-[0-[3-[4-(phenylmethylamino)butylamino]propyl]-oxime] (Compound 13);

^{13}C -NMR (50 MHz, CDCl_3): δ (ppm): 175.22; 171.45; 140.35; 128.39;
128.13; 126.90; 102.98; 96.26; 53.94;

Erythromycin A (E)-9-[0-[6-[N-isopropyl-2-(2-phenylethylamino)ethyl]-

5 aminohexylloxime] (Compound 14)

m.p. 93-95°C

Mass (C.I.) (M+H) $^+$ =1038

^{13}C -NMR (50 MHz, CDCl_3): δ (ppm): 174.92; 170.96; 139.71; 128.42;
128.10; 125.79; 102.62; 95.94; 50.93;

10 Erythromycin A (E)-9-[0-[6-[4-(N-isopropyl-phenylmethylamino)butyl]-
aminohexylloxime] (Compound 15)

m.p. 78-80°C

Mass (C.I.) (M+H) $^+$ =1052

^{13}C -NMR (50 MHz, CDCl_3): δ (ppm): 175.47; 171.30; 140.95; 128.61;
15 128.02; 126.70; 116.87; 102.94; 53.94;

Erythromycin A (E)-9-[0-[2-[6-[(4-fluorophenyl)methylamino]hexylami-
no]ethylloxime] (Compound 16)

Mass (C.I.) (M+H) $^+$ =999.5

^{13}C -NMR (50 MHz, CDCl_3): δ (ppm): 161.88; 136.06; 129.65; 115.14;

20 Erythromycin A (E)-9-[0-[2-[6-[(4-methoxyphenyl)methylamino]hexyl-
amino]ethylloxime] (Compound 17)

Mass (C.I.) (M+H) $^+$ =1011

^{13}C -NMR (50 MHz, CDCl_3): δ (ppm): 158.57; 132.58; 129.31; 113.76;

Erythromycin A (E)-9-[0-[2-[6-[(3,4-methylenedioxyphenyl)methylami-
25 no]hexylamino]ethylloxime] (Compound 18)

Mass (C.I.) (M+H) $^+$ =1025

^{13}C -NMR (50 MHz, CDCl_3): δ (ppm): 147.65; 146.44; 134.39; 121.18;
108.66, 108.06;

Erythromycin A (E)-9-[0-[2-[6-[(3-trifluoromethylphenyl)methylami-
30 no]hexylamino]ethylloxime] (Compound 19)

Mass (C.I.) (M+H)⁺=1050

¹³C-NMR (50 MHz, CDCl₃): δ (ppm): 141.57; 131.40; 130.63; 128.76;
124.22; 124.72; 123.56;

5 Erythromycin A (E)-9-[0-[2-[6-[(4-methylsulphonylphenyl)methylamino]hexylamino]ethyl]oxime] (Compound 20)

Mass (C.I.) (M+H)⁺=1059

¹³C-NMR (50 MHz, CDCl₃): δ (ppm): 147.23; 138.97; 128.79; 127.49.

Example 13

10 Preparation of N-benzyloxycarbonyl-6-aminohexanol

Benzyl chloroformate (50% in toluene; 84.8 ml; 0.256 moles) in ethyl acetate (171 ml) and a solution of sodium hydroxide 1N (256 ml) were gradually and contemporaneously added to a mixture of 6-aminohexanol (25 g; 0.21 moles) in ethyl acetate (250 ml) and water (200 ml).

15 kept under stirring at 0°C.

The reaction mixture (pH 9) was brought at room temperature and kept under stirring for 5 hours.

After separation of the phases, the aqueous phase was washed with ethyl acetate (200 ml).

20 The collected organic phases were then washed with a saturated solution of sodium chloride (150 ml), dried on sodium sulphate and evaporated to dryness.

The residue was collected with ethyl ether (300 ml) and the formed precipitate was filtered and dried under vacuum at 50°C, affording

25 thus N-benzyloxycarbonyl-6-aminohexanol (44.5 g).

m.p. 80-82°C.

Example 14

Preparation of N-benzyloxycarbonyl-6-amino-hexanal

A solution of potassium bromide (1.89 g; 16 mmoles) in water (31 ml) was added to a solution of N-benzyloxycarbonyl-6-aminohexanol (40 g;

- 39 -

0.159 moles), prepared as described in example 13, in methylene chloride (600 ml) containing the free radical 2,2,6,6-tetramethyl-piperidinoxy (TEMPO) (0.248 g; 1.6 mmols).

- 5 A solution of sodium hypochlorite (215 ml), prepared by mixing a solution of sodium hypochlorite at 7% (240 ml) with sodium bicarbonate (4.22 g) and hydrochloric acid at 5% (5 ml), in order to reach pH 8.7, was gradually added to the reaction mixture, kept under stirring at a temperature of 10°C.
- 10 At the end of the addition, after separation of the phases, the organic phase was washed with methylene chloride (2x200 ml), dried on sodium sulphate and evaporated to dryness.
- N-benzyloxycarbonyl-6-amino-hexanal (39.45 g) was thus obtained as an oil.

- 15 TLC (ethyl acetate:hexane=1:1) Rf=0.41.

Example 15

Preparation of 2-[6-(benzyloxycarbonylamino)hexylamino]ethanol

- A mixture constituted by N-benzyloxycarbonyl-6-amino-hexanal (35 g; 0.14 moles) and 2-aminoethanol (51.3 g; 0.84 moles) in ethanol (250
- 20 ml), in the presence of molecular sieves (3 Å), was kept under stirring at room temperature for two hours.

The reaction mixture was then filtered on celite and sodium boron hydride (6.33 g; 0.168 moles) was added to the resultant solution.

- After 4 hours under stirring at room temperature, the reaction
- 25 solvent was evaporated under vacuum and the residue was collected with water (500 ml) and ethyl acetate (500 ml).

After separation of the phases, the aqueous phase was further extracted with ethyl acetate (200 ml).

- The collected organic phases were washed with a saturated solution
- 30 of sodium chloride (250 ml), dried on sodium sulphate and evaporated

- 40 -

to dryness, obtaining thus 2-[6-(benzyloxycarbonylamino)hexylamino]ethanol (38.36 g).

TLC (ethyl acetate:methanol:ammonia=10:2:1) Rf=0.4.

5

Example 16

Preparation of 2-[N-benzyloxycarbonyl-6-(benzyloxycarbonylamino)hexylamino]ethanol

By working analogously to what described in example 9 and by using 2-[6-(benzyloxycarbonylamino)hexylamino]ethanol (38.3 g; 0.13 moles), prepared as described in example 15, 2-[N-benzyloxycarbonyl-6-(benzyloxycarbonylamino)hexylamino]ethanol was obtained as an oil. TLC (ethyl acetate:hexane=65:35) Rf=0.45

Example 17

Preparation of 2-[N-benzyloxycarbonyl-6-(benzyloxycarbonylamino)hexylamino]ethyl-methanesulphonate

By working analogously to what described in example 10 and by using 2-[N-benzyloxycarbonyl-6-(benzyloxycarbonylamino)hexylamino]ethanol (20 g; 47.8 mmols), prepared as described in example 16, 2-[N-benzyloxycarbonyl-6-(benzyloxycarbonylamino)hexylamino]ethyl-methanesulphonate (24.35 g) was obtained as an oil, used as such in the subsequent reactions.

Example 18

Preparation of Erythromycin A (E)-9-[0-[2-[N-benzyloxycarbonyl-6-(benzyloxycarbonylamino)hexylamino]ethyl]oxime]

By working analogously to what described in example 11 and by using 2-[N-benzyloxycarbonyl-6-(benzyloxycarbonylamino)hexylamino]ethyl-methanesulphonate (24.25 g; 47.8 mmols), prepared as described in example 17, after silica gel chromatography (eluent methylene chloride:methanol:ammonia=95:5:0.5), Erythromycin A (E)-9-[0-[2-[N-benzyloxycarbonyl-6-(benzyloxycarbonylamino)hexylamino]ethyl]oxime]

30

- 41 -

(36.1 g) was obtained.

TLC (methylene chloride:methanol:ammonia=85:15:1.5) Rf=0.5

¹H-NMR (200 MHz, CDCl₃): δ (ppm): 7.39-7.22 (m, 10H, aromatics);
5 5.14-5.05 (m, 4H, 2 CH₂Ph); 3.29 (s, 3H, OCH₃); 2.25 (s, 6H, 2
NCH₃); 0.80 (t, 3H, CH₂CH₃).

Example 19

Preparation of Erythromycin A (E)-9-[0-[2-(6-amino-hexylamino)ethyl- ylloxime]

- 10 By working analogously to what described in example 12 and by using
Erythromycin A (E)-9-[0-[2-[N-benzyloxycarbonyl-6-(benzyloxycarbon-
ylamino)hexylamino]ethylloxime], prepared as described in example
18, after silica gel chromatography (eluent methylene chloride:
methanol:ammonia=85:15:1.5), Erythromycin A (E)-9-[0-[2-(6-amino-
15 -hexylamino)ethyl]oxime] was obtained.

TLC (methylene chloride:methanol:ammonia=85:15:1.5) Rf=0.2

¹³C-NMR (50 MHz, CDCl₃): δ (ppm): 175.18; 171.26; 102.96; 96.28.

Example 20

- 20 Preparation of Erythromycin A (E)-9-[0-[2-[6-[(2-trifluoromethyl-
phenyl)methylaminohexylamino]ethylloxime] (Compound 21)

2-Trifluoromethylbenzaldehyde (0.4 g) and molecular sieves (4.5 g; 3
A) were added to a solution of Erythromycin A (E)-9-[0-[2-(6-amino-
hexylamino)ethylloxime] (2 g; 2.24 mmol), prepared as described in
example 19, in ethanol (50 ml), kept under stirring at room temper-
25 ature.

After 2 hours, the molecular sieves were filtered off and palladium
on charcoal at 10% (0.2 g) was added to the resultant solution.

The reaction mixture was placed into a Parr hydrogenator which was
loaded with hydrogen (1 bar).

- 30 After one hour, ended the hydrogenation reaction, the catalyst was

filtered off and the solvent was evaporated.

The residue was purified by silica gel chromatography (eluent methylene chloride:methanol:ammonia=95:5:0.5), thus obtaining Erythromycin A (E)-9-[0-[2-[6-[(2-trifluoromethylphenyl)methylamino]hexylamino]ethylloxime] (2 g).

Mass (C.I.) (M+H)⁺=1050

¹³C-NMR (50 MHz, CDCl₃): δ (ppm): 139.14; 131.88; 130.38; 127.58; 126.81; 125.82.

10 By working analogously the following compounds were prepared:

Erythromycin A (E)-9-[0-[2-[6-(3-pyridylmethylamino)hexylamino]ethylloxime] (Compound 22)

Mass (C.I.) (M+H)⁺=982

¹³C-NMR (50 MHz, CDCl₃): δ (ppm): 149.66; 148.39; 135.81; 123.40;

15 Erythromycin A (E)-9-[0-[2-[6-(4-trifluoromethylphenyl)methylamino]hexylamino]ethylloxime] (Compound 23)

Mass (C.I.) (M+H)⁺=1050

¹³C-NMR (50 MHz, CDCl₃): δ (ppm): 144.73; 128.23; 125.25; 124.26;

20 Erythromycin A (E)-9-[0-[2-[6-(2-hydroxyphenyl)methylamino]hexylamino]ethylloxime] (Compound 24)

Mass (C.I.) (M+H)⁺=997

¹³C-NMR (50 MHz, CDCl₃): δ (ppm): 158.37; 128.60; 128.19; 122.54; 118.88; 116.32;

25 Erythromycin A (E)-9-[0-[2-[6-(3-hydroxyphenyl)methylamino]hexylamino]ethylloxime] (Compound 25)

Mass (C.I.) (M+H)⁺=997

¹³C-NMR (50 MHz, CDCl₃): δ (ppm): 157.28; 140.46; 129.56; 119.70; 115.55; 114.89;

30 Erythromycin A (E)-9-[0-[2-[6-(4-n.butoxyphenyl)methylamino]hexylamino]ethylloxime] (Compound 26)

Mass (C.I.) (M+H)⁺=1053

¹³C-NMR (50 MHz, CDCl₃): δ (ppm): 158.27; 131.65; 129.40; 114.40;

Erythromycin A (E)-9-[0-[2-[6-[(3-phenoxyphenyl)methylamino]hexyl-

5 aminoethylloxime] (Compound 27)

Mass (C.I.) (M+H)⁺=1073

¹³C-NMR (50 MHz, CDCl₃): δ (ppm): 157.32; 157.28; 142.69; 129.72;

129.64; 123.16; 122.91; 118.84; 118.52; 117.29;

Erythromycin A (E)-9-[0-[2-[6-[(4-hydroxyphenyl)methylamino]hexyl-

10 aminoethylloxime] (Compound 28)

Mass (C.I.) (M+H)⁺=997

¹³C-NMR (50 MHz, CDCl₃): δ (ppm): 156.49; 130.00; 128.87; 115.88;

Erythromycin A (E)-9-[0-[2-[6-[(4-phenoxyphenyl)methylamino]hexyl-

aminoethylloxime] (Compound 29)

15 Mass (C.I.) (M+H)⁺=1073

¹³C-NMR (50 MHz, CDCl₃): δ (ppm): 157.43; 156.05; 135.43; 129.69;

129.49; 123.07; 118.92; 118.72; 118.67;

Erythromycin A (E)-9-[0-[2-[6-[(biphenyl-4-yl)methylamino]hexyl-

aminoethylloxime] (Compound 30)

20 Mass (C.I.) (M+H)⁺=1057

¹³C-NMR (50 MHz, CDCl₃): δ (ppm): 140.94; 139.86; 139.40; 128.74;

128.58; 127.13; 127.03;

Erythromycin A (E)-9-[0-[2-[6-[(2-furylmethylamino)hexylamino]-

ethylloxime] (Compound 31)

25 Mass (C.I.) (M+H)⁺=971

¹³C-NMR (50 MHz, CDCl₃): δ (ppm): 153.92; 141.73; 110.08; 106.81.

Example 21

Preparation of Erythromycin A (E)-9-[0-[2-[6-[(3,5-dichloro-2-hydro-

xyphenyl)methylamino]hexylaminoethylloxime] (Compound 32)

Molecular sieves (6 g; 3 Å) and 3,5-dichloro-2-hydroxybenzaldehyde

- 44 -

(0.535 g; 2.8 mmoles) were added to a solution of Erythromycin A (E)-9-[0-[2-(6-amino-hexylamino)ethyl]oxime] (2.5 g; 2.8 mmoles), prepared as described in example 19, in anhydrous ethanol (100 ml).

- 5 The reaction mixture was kept under stirring at room temperature and, after 2 hours, the molecular sieves were filtered off and sodium boron hydride (0.106 g; 2.89 mmoles) was added, portionwise, to the resultant solution.

After 3 hours under stirring, the solvent was evaporated at reduced
10 pressure and the residue was purified by silica gel chromatography (eluent methylene chloride:methanol:ammonia=85:15:1.5) obtaining thus Erythromycin A (E)-9-[0-[2-[6-[(3,5-dichloro-2-hydroxyphenyl)-methylamino]hexylamino]ethyl]oxime] (2.2 g).

Mass (C.I.) (M+H)⁺=1066

- 15 ¹³C-NMR (50 MHz, CDCl₃): δ (ppm): 153.43; 128.43; 126.42; 124.41; 122.91; 121.61.

By working analogously the following compounds were prepared:

Erythromycin A (E)-9-[0-[2-[6-[(2-nitrophenyl)methylamino]hexylamino]ethyl]oxime] (Compound 33)

- 20 Mass (C.I.) (M+H)⁺=1027

¹³C-NMR (50 MHz, CDCl₃): δ (ppm): 149.14; 135.79; 133.13; 131.26; 127.87; 124.70;

Erythromycin A (E)-9-[0-[2-[6-[(3-nitrophenyl)methylamino]hexylamino]ethyl]oxime] (Compound 34)

- 25 Mass (C.I.) (M+H)⁺=1027

¹³C-NMR (50 MHz, CDCl₃): δ (ppm): 148.37; 142.87; 134.17; 129.22; 122.81; 121.96;

Erythromycin A (E)-9-[0-[2-[6-[(4-nitrophenyl)methylamino]hexylamino]ethyl]oxime] (Compound 35)

- 30 Mass (C.I.) (M+H)⁺=1027

^{13}C -NMR (50 MHz, CDCl_3): δ (ppm): 148.41; 147.00; 128.59; 123.60;

Erythromycin A (E)-9-[0-[2-[6-[(4-hydroxy-3-nitrophenyl)methylamino]hexylamino]ethyl]oxime] (Compound 36)

5 Mass (C.I.) $(\text{M}+\text{H})^+=1043$

^{13}C -NMR (50 MHz, CDCl_3): δ (ppm): 157.29; 137.40; 134.05; 128.01; 125.23; 121.70;

Erythromycin A (E)-9-[0-[2-[6-[(3-hydroxy-4-nitrophenyl)methylamino]hexylamino]ethyl]oxime] (Compound 37)

10 Mass (C.I.) $(\text{M}+\text{H})^+=1043$

^{13}C -NMR (50 MHz, CDCl_3): δ (ppm): 155.50; 151.98; 132.51; 125.13; 119.67; 118.59;

Example 22

Preparation of Erythromycin A (E)-9-[0-[2-[N-methyl-6-(N'-methyl-N'-phenylmethylamino)hexylamino]ethyl]oxime] (Compound 38)

15 An aqueous solution of formaldehyde at 37% (2 ml; 26.6 mmol) and palladium on charcoal at 10% (0.82 g) were added, in this order, to a solution of Erythromycin A (E)-9-[0-[2-[6-(phenylmethylamino)hexylamino]ethyl]oxime] (2 g; 2 mmol), prepared as described in
20 example 12, in a mixture ethanol:water=1:1 (20 ml) kept under stirring at room temperature.

The reaction mixture was placed into a Parr hydrogenator loaded with hydrogen (1 bar).

25 After 2 hours, the reaction mixture was filtered to eliminate the catalyst and the resultant solution was evaporated to dryness.

The obtained residue was purified by silica gel chromatography (eluent methylene chloride:methanol:ammonia=90:10:1) affording Erythromycin A (E)-9-[0-[2-[N-methyl-6-(N'-methyl-N'-phenylmethylamino)hexylamino]ethyl]oxime] (1.8 g).

30 Mass (C.I.) $(\text{M}+\text{H})^+=1009$

- 46 -

^{13}C -NMR (50 MHz, CDCl_3): δ (ppm): 139.20; 129.04; 128.17; 126.86.

By working analogously the following compound was prepared:

Erythromycin A (E)-9-[0-[2-[N-methyl-6-[N'-methyl-N'-(4-trifluoro-
5 methylphenyl)methylamino]hexylamino]ethylloxime] (Compound 39)

Mass (C.I.) $(\text{M}+\text{H})^+=1078$

^{13}C -NMR (50 MHz, CDCl_3): δ (ppm): 143.65; 129.12; 129.03; 125.10;
124.29.

Example 23

10 Pharmacologic activity

a) In vitro antibacterial activity

The determination of the minimum inhibiting concentrations (MIC), with respect to Gram-positive and Gram-negative bacteria was carried out through the micromethod of gradual broth dilution in
15 double series [National Committee for Clinical Laboratory Standards, 1990; Methods for dilution antimicrobial susceptibility tests for bacteria that grow aerobically; Approved standards M7-A2-NCCLS, Villanova, Pa.], by using Mueller Hinton Broth (MHB) as a culture medium.

20 In the case of exigent bacteria, horse serum at 5% (Streptococcus pneumoniae and Streptococcus pyogenes) was added to the medium. Roxithromycin and Clarithromycin [The Merck Index, XI Ed., No. 8253 and 2340, respectively] were used as reference macrolides. MIC, expressed as ($\mu\text{g}/\text{ml}$), were determined after incubation of
25 the microplates at 37°C for 18 hours, evaluating the lowest antibiotic concentration enabling the inhibition of bacterial development.

b) In vivo antibacterial activity

The therapeutical effectiveness, expressed as average protecting
30 dose (PD_{50}), of the considered compounds of formula (I) was

evaluated by experimental pulmonary infection induced in mouse by *Streptococcus pyogenes* C 203.

Charles River albin mice (strain CD 1), of body weight comprised
5 between 23-35 g, stabulated in groups of 6 for cage and
normo-fed with standard diet and water ad libitum were used.

A suspension of *S. pyogenes* C 203 (corresponding about to 10⁸
UFC) in tryptose broth (0.05 ml) was intranasally administered to
each mouse anesthetized with a mixture of ethyl ether and chlo-
10 roform.

The compounds under examination were intraperitoneally adminis-
tered as a single dose, in 0.2% Tween suspension, 24 hours before
or 1 hour after the infection.

Mice mortality observation was prolonged up to 10 days from the
15 infection.

PD₅₀ calculation, expressed as (mg/Kg), was carried out through
the analysis of probit.

For some representative compounds of formula (I) the in vitro anti-
bacterial activity against Gram-positive microorganisms (table 1)
20 and Gram-negative microorganisms (table 2), and the in vivo anti-
bacterial activity values (table 3), are reported as follows.

Table 1

In vitro antibacterial activity, expressed as minimum inhibiting
concentration MIC (μ g/ml), of the compounds 2, 4-12, 16-19, 21,
25 23-38 and of the reference compounds Roxithromycin and Clarithromy-
cin, against Gram-positive microorganisms such as *Streptococcus*
pneumoniae BS 3, *Streptococcus pneumoniae* BS 4, *Streptococcus*
pyogenes A 26, *Streptococcus pyogenes* C 203, *Enterococcus faecalis*
ATCC 29212 and *Staphylococcus aureus* PV 14.

- 48 -

		MIC ($\mu\text{g/ml}$)					
compound		S. pneumoniae BS 3	S. pneumoniae BS 4	S. pyogenes A 26	S. pyogenes C 203	E. faecalis ATCC 29212	S. aureus PV 14
5	2	0.0156	0.0312	0.0156	0.0312	8	1
	4	0.0156	0.0156	0.0156	0.0039	4	0.25
	5	0.0156	0.0312	0.0312	0.0156	4	1
	6	0.0156	0.0312	0.0312	0.0156	8	0.25
	7	0.0625	0.0625	0.0312	0.0156	4	0.5
10	8	0.25	0.5	0.0312	0.0078	8	0.5
	9	0.0078	0.0078	0.0078	0.0039	4	0.25
	10	0.0156	0.0156	0.0078	0.0039	4	0.5
	11	0.25	0.5	0.0625	0.0312	16	0.5
	12	0.25	0.5	0.0625	0.0312	16	0.25
15	16	0.0156	0.0156	0.0156	0.0039	4	0.25
	17	0.0312	0.0625	0.0625	0.0039	4	0.25
	18	0.0156	0.0312	0.0312	0.0078	2	0.25
	19	0.0156	0.0078	0.0156	0.0156	2	1
	21	0.0078	0.0039	0.0078	0.0156	2	1
20	23	0.0156	0.0078	0.0156	0.0156	2	1
	24	0.0078	0.0156	0.0156	0.0039	4	0.25
	25	0.25	0.25	0.125	0.0312	8	0.125
	26	0.0156	0.0156	0.0312	0.0078	1	0.25
	27	0.0078	0.0078	0.0156	0.0078	1	0.5
25	28	0.25	0.25	0.25	0.125	16	0.125
	29	0.0078	0.0156	0.0312	0.0039	1	0.5

- 49 -

compound	MIC ($\mu\text{g/ml}$)					
	S. pneumoniae BS 3	S. pneumoniae BS 4	S. pyogenes A 26	S. pyogenes C 203	E. faecalis ATCC 29212	S. aureus PV 14
5						
30	0.0156	0.0156	0.0156	0.0078	1	0.5
31	0.0078	0.0078	0.0078	0.0039	4	0.5
32	0.0625	0.0312	0.0625	0.0156	2	0.5
33	0.0039	0.0039	0.0039	0.00097	2	0.5
34	0.0019	0.0039	0.0078	0.0039	1	0.25
10						
35	0.0039	0.0039	0.0078	0.0019	0.5	0.25
36	0.0625	0.0625	0.0625	0.0078	16	0.5
37	0.0312	0.0312	0.0312	0.0039	8	0.5
38	0.0078	0.0039	0.0156	0.0078	8	0.5
Roxithromycin	0.0312	0.0625	0.0625	0.0625	4	1
15						
Clarithromycin	0.0078	0.0156	0.0078	0.0078	1	0.25

The above reported data clearly indicate that the compounds of formula (I), object of the present invention, are endowed with an antibacterial activity substantially comparable to that of Clarithromycin and Roxithromycin, with respect to Gram-positive microorganisms.

Table 2

In vitro antibacterial activity, expressed as minimum inhibiting concentration MIC ($\mu\text{g/ml}$), of the compounds 2, 4-12, 16-19, 21, 23-38 and of the reference compounds Roxithromycin and Clarithromycin, against Gram-negative microorganisms such as Escherichia coli ATCC 25922 and Klebsiella pneumoniae ZC 2.

- 50 -

	Compound	MIC ($\mu\text{g/ml}$)	
		E. coli ATCC 25922	K. pneumoniae ZC 2
5	2	16	64
	4	4	16
	5	8	32
	6	4	16
	7	4	16
10	8	4	16
	9	4	16
	10	4	16
	11	8	16
	12	8	32
15	16	2	8
	17	4	16
	18	2	16
	19	2	8
	21	4	16
20	23	1	4
	24	4	8
	25	8	16
	26	1	2
	27	1	2
25	28	16	32
	29	1	2
	30	1	2
	31	4	16

- 51 -

	Compound	MIC ($\mu\text{g/ml}$)	
		E. coli ATCC 25922	K. pneumoniae ZC 2
5	32	4	16
	33	4	16
	34	1	8
	35	1	4
	36	8	32
10	37	4	16
	38	8	32
	Roxithromycin	28	256
	Clarithromycin	64	128

15 The antibacterial activity of the compounds of formula (I) against Gram-negative microorganisms such as *Escherichia coli* and *Klebsiella pneumoniae* resulted to be markedly higher than that of both reference compounds.

Table 3

20 In vivo antibacterial activity, expressed as average protecting dose PD_{50} (mg/Kg), 24 hours before and 1 hour after the experimental pulmonary infection induced in mouse by *Streptococcus pyogenes* C 203, of the compounds 4, 10, 16-19, 21, 23, 26-27, 29-30, 33-35 and 38 and of the reference compounds Roxithromycin and Clarithromycin.

25

30


- 52 -

	Compound	PD ₅₀ (mg/Kg) Pulmonary infection (S. pyogenes C 203)	
		1 hour after infection	24 hours before infection
5	4	0.9	4.78
	10	2.36	5.8
	16	3.6	10.21
	17	1.17	8.16
	18	1.32	2.23
10	19	1.95	4.84
	21	0.82	5.95
	23	1.46	3.49
	26	6.11	6.11
	27	15.6	15.6
15	29	7.65	6.1
	30	19.3	19.3
	33	1.61	6.11
	34	1.88	6.8
	35	3.00	6.0
20	38	11.8	11.8
	Roxithromycin	0.9	>25
	Clarithromycin	3.25	>50

The compounds of formula (I) resulted to be active in vivo and their activity profile indicates that said compounds present a duration of action and a half-life of tissue elimination significantly higher than those of both reference compounds.

Claims

1) A compound of formula



wherein

A is a phenyl group or a heterocycle with 5 or 6 members containing 1 or more heteroatoms selected among nitrogen, oxygen and sulphur, optionally substituted with from 1 to 3 groups, the same or different, selected among straight or branched C₁-C₄ alkyl or alkoxy groups, C₁-C₂ alkylenedioxy groups, C₁-C₄ alkylsulphonyl groups, phenyl, phenoxy, hydroxy, carboxy, nitro, halogen and trifluoromethyl groups;


R₁ and R₂, the same or different, represent a hydrogen atom or a straight or branched C₁-C₄ alkyl group;

n is 1 or 2;

m is an integer comprised between 1 and 8;

r is an integer comprised between 2 and 6;

M represents a group of formula



wherein

R₁ is a hydrogen atom or a methyl group;

and its pharmaceutically acceptable salts.

5 2) A compound according to claim 1 having E configuration.

3) A compound according to claim 1 wherein A represents a phenyl group or a heterocycle selected between pyridine and furan, optionally substituted with from 1 to 3 groups selected among hydroxy, methoxy, methylenedioxy, n.butoxy, phenoxy, phenyl, methylsulphonyl, nitro, halogen and trifluoromethyl groups; R₁ and
10 R₂, the same each other, represent a hydrogen atom or a methyl group; R₃ represents a hydrogen atom.

4) A compound according to claim 1 wherein A represents a phenyl group optionally substituted with a group selected among phenoxy, nitro and trifluoromethyl; R₁ and R₂, the same each other, represent
15 a hydrogen atom or a methyl group; n is equal to 1; m is equal to 6; p is equal to 2; R₃ represents a hydrogen atom.

5) A pharmaceutical composition containing a therapeutically effective amount of one or more compounds of formula (I) in admixture with a pharmaceutically acceptable carrier.

20 6) A method for the treatment of infectious diseases in human or veterinary therapy consisting in administering a therapeutically effective amount of a compound according to claim 1.

7) A method according to claim 6 for the treatment of malarian diseases.

25

INTERNATIONAL SEARCH REPORT

International Application No.

PCT/EP 95/04815

A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 C07H17/08 A61K31/70

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07H A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	EP,A,0 487 411 (ROUSSEL UCLAF) 27 May 1992 see page 37 - page 38; examples ---	1-6
X	EP,A,0 033 255 (ROUSSEL UCLAF) 5 August 1981 cited in the application see page 1 - page 7 ---	1-6
A	US,A,4 740 502 (HANNICK STEVEN M ET AL) 26 April 1988 see the whole document ---	1,5,6
A	EP,A,0 422 843 (TAISHO PHARMA CO LTD) 17 April 1991 see the whole document ---	1,5,6
-/--		

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

A document member of the same patent family

Date of the actual completion of the international search

23 April 1996

Date of mailing of the international search report

17.05.96

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl,
Fax (+ 31-70) 340-3016

Authorized officer

Moreno, C

INTERNATIONAL SEARCH REPORT

International Application No
PCT/EP 95/04815

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>JOURNAL OF ANTIBIOTICS, vol. 44, no. 3, March 1991, TOKYO JP, pages 313-330, XP000567789 J. GASC ET AL: "New ether oxime derivatives of erythromycin A. A structure-activity relationship study." see the whole document -----</p>	1,5,6

INTERNATIONAL SEARCH REPORT

International application No.

PCT/EP 95/04815

Box I Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

Remark: Although claims 6 and 7 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of Item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 95/04815

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP-A-0487411	27-05-92	FR-A- 2669337	22-05-92
		FR-A- 2677025	04-12-92
		FR-A- 2680790	05-03-93
		AT-T- 133683	15-02-96
		AU-B- 640290	19-08-93
		AU-B- 8798691	28-05-92
		CA-A- 2055912	21-05-92
		CN-A- 1065069	07-10-92
		DE-D- 69116815	14-03-96
		JP-A- 4290893	15-10-92
		NZ-A- 240684	26-08-94
		OA-A- 9523	15-11-92
		PL-B- 167448	30-09-95
		US-A- 5444051	22-08-95
EP-A-0033255	05-08-81	FR-A- 2473525	17-07-81
		AT-T- 9161	15-09-84
		AU-B- 537247	14-06-84
		AU-B- 6613181	16-07-81
		CA-A- 1162535	21-02-84
		JP-C- 1576270	24-08-90
		JP-A- 56100799	12-08-81
		JP-B- 62037639	13-08-87
		LU-A- 88283	09-09-94
		OA-A- 6719	30-06-82
		US-A- 4349545	14-09-82
US-A-4740502	26-04-88	NONE	
EP-A-0422843	17-04-91	AT-T- 118503	15-03-95
		CA-A- 2027074	08-04-91
		DE-D- 69016912	23-03-95
		DE-T- 69016912	06-07-95
		ES-T- 2071036	16-06-95
		JP-A- 3204893	06-09-91
		US-A- 5302705	12-04-94